

# ORGANIC AMINO ACID FERTILIZERS

**ROK**  **AKTIV**

**ROK**  **OLAN**

**ROK**  **HUMIN**

## FOR HEALTHIER AND SELF-SUFFICIENT WORLD

Scan QR code  
for online version



**ROK**  **SAN**<sup>®</sup>

 **SLOVAK  
PAVILION**

# PROLOGUE

In recent uncertain times, with the borders closed, the need for food self-sufficiency has been more important than ever. Food self-sufficiency comes hand in hand with food safety. What you are holding now in your hands is a roadmap to food self-sufficiency and safety of the planet Earth.

Our liquid amino acid fertilizers ROKOAKTIV, ROKOLAN and ROKOHUMIN are designed to be used in organic farming and to assure high yield and product quality. Fruits are sweeter and bigger, but more importantly, healthier. Amino acids are essential in crop production in regards to their effect on immunity of the plants itself and quality of the yield.

In the world where the prices of chemical fertilizers are on the rise, we can offer stability for farmers and even saving on fertilization costs by 30 %.

By applying our complex nutritional technology for treating seeds, soil and plant (leaves), we strengthen the human body's immunity which is important especially in these difficult times when human organisms are being attacked by various viral diseases. Furthermore, fertilizers contain carbon that is so much needed for soil regeneration and revitalization. Healthy soil is a key for CO2 sequestration and helps fight global warming. Strong plants are able to withstand extreme weather conditions such as droughts and floods. The technological process is patented in the USA, Russia, Ukraine and China. Fertilizer is being sold in Europe, Africa, Russia and Latin America.

Our goal is to start producing locally in every country where there is an abundance of raw materials, by using local workforce. We believe that our mission is to make the world better and healthier. Countries will become self-sufficient in production of organic fertilizers, which is a step towards food security.

We believe that our ideas and technologies are beneficial for whole planet. Combination of organic fertilizers with scientific methods of water management can make world greener and healthier place.



EXPO  
2020  
DUBAI  
UAE



SLOVAK  
PAVILION

OFFICIAL PARTICIPANT

# WE ARE RENEWING LIFE IN THE SOIL AND HELPING TO SEQUESTER CARBON FROM THE ATMOSPHERE

Many of our clients have understood that their yields are in many cases higher than in the past, when they used industrial fertilizers and they return to us with trust.

Not to mention that **the fertilizing cost per 1 ha is 30% lower compared to industrial fertilizers.**

Clients even perceive that life in their soil is being renewed.

Healthy soil sequesters and hold CO<sub>2</sub> thus preventing it from escaping into our atmosphere. **The higher the organic matter in the soil, the higher is the soil's ability to sequester CO<sub>2</sub>.**

Deforestation, loss of soil nutrients, loss of biodiversity, excessive agricultural practices (compaction, excessive chemical fertilizing), erosion, waterlogging and urbanization cause our soil to emit greenhouse gases and contribute to climate change and global warming.

**Our fertilizers contain carbon and amino acids that enrich and revitalize soil microorganisms that help to retain carbon in the soil and accelerate the plant's growth. Carbon rich soil absorbs more water which can be beneficial for the plant during drought as well as during floods.**

**Sustainable farming practices help to keep soil healthy and to fight climate change.**



The picture shows the pre-sowing soil preparation with Rokolan 50 lit./ha




**WE ARE RENEWING LIFE IN THE SOIL**

# EFFICIENCY OF FERTILIZER



Field tests on Wheat 2019-2020\*



\* Conducted by NATIONAL AGRICULTURAL AND FOOD CENTRE - Agroecology Research Institute in Michalovce, Slovak Republic



## Enhancement of soil quality and fertility

Thanks to humic and fulvic acids we are able to increase the fertility and quality of soil by multiplication of living microorganisms and absorption of heavy metals and other pollutants



## Support of plant immunity system

Essential amino acids support immunity system and protect plants from stress factors such as drought, frost or diseases



## Increase of yield quantity and quality

Balanced organic nutrition offered by ROKOSAN products has positive effect on formation of protein and sugar in crops along with shortened time of ripening



## Our products are environmentally friendly

Thanks to organic nature of our fertilizers, they are not polluting the groundwater nor are toxic to the animals or bees

# FINANCIAL BENEFITS OF ROKOSAN TECHNOLOGY

Our **ROKOLAN + ROKOAKTIV + ROKOHUMIN** COMPETE WITH **CHEMICAL FERTILIZERS** in terms of price and results



Our **FERTILIZERS** are **HIGH QUALITY PRODUCTS** that are harmless to soil, while ensuring **HEALTHY NUTRITION AND IMMUNITY**



While the prices of chemical fertilizers are rising every day, which is causing the subsequent rise in food prices, the **PRICE AND QUALITY** of our products **ROKOLAN, ROKOAKTIV and ROKOHUMIN** REMAIN **UNCHANGED**



## PRICE COMPARISON PER HECTARE



\* Calculated for wheat cultivation \*\* Based on prices from November 2021 in European Markets

# WATER MANAGEMENT SOLUTION



WATER HOLISTIC

We are closely cooperating with **WATERHOLISTIC** company.

**Together, we can offer comprehensive solution to the clients that not only need to revitalize their soil but also bring water and biodiversity back to their land.**

Humanity, through deforestation, agriculture and urbanization accelerates the runoff of rainwater and drains the land. Heat Islands or heat domes islands" humans create cause breakdowns in the stable water cycle, drive climate extremes, more frequent flooding, longer droughts, extreme heat, forest fires, drop in the groundwater reserves, and decrease in soil fertility and biodiversity.

However, the impact can go in both directions. **The part of climate change, which is the result of human draining of land, can be stopped. The renewal of a healthy water cycle can be achieved by a comprehensive program of rainwater harvesting, conservation, infiltration, and evaporation in each watershed.**

Founders of WATERHOLISTIC have more than 30 years of experience with water retention measures, land management and restoration of small water cycles.

Experiences of founders involve:

- **Alternative Blue Water Management plan for Slovakia (1993 - 94)**
- **Stop Drying of Slovakia proposal or policy (1998 – 2000)**
- **Sustainable city water management Kosice Water Protocol (2005)**
- **Several programs of alternative water resources creation and increasing water retention in natural ecosystems 1995 - 2008)**
- **Educational activities like School of Regional Development (2007 - 2008) and People's University of Water (2007 - 2008)**
- **Management of government program of landscape restoration (2011 – 2012)**
- **Green restoration program for the Kosice region of Slovakia: Landscape and watershed recovery (2021-2030)**





# PILLARS OF A QUALITY YIELD



**ROKOAKTIV**  
Germination  
activator

+



**ROKOLAN**  
Soil  
activator

+



**ROKOHUMIN**  
Leaf  
nutrition



These products can be used standalone in the technological processes, but excellent results in soil recovery and fertility growth are achieved by **combining all three fertilizers.**

Amino acid fertilizers containing **18 basic amino acids.**  
In addition, Rokoaktiv and Rokohumin contain **humic acids.**





Rwanda



Slovakia



Estonia



Poland



Serbia



Ivory Coast



California



Spain



Ukraine



Canada



Tanzania



Colombia



Cuba



Hungaria



Kenya



Czech Republic



Belarus



Moldova



Panama

# LEAF NUTRITION



The advantage of foliar nutrition lies in that it provides enough varied nutrients, which are nearly essential, to the plants. It fixed the deficit most effectively and quickly, or if intake through the roots is for some reason limited (eg pH of the environment). Leaf nutrition is quickly taken up by the plant and its effect is therefore visible the fastest.

Rokohumin is a nutritional cocktail of amino acids, peptides, humic acids, macro and micronutrients – that’s why it’s the right choice for plants.

### Rokohumin can be used:

1. During each phase of the growth
2. In the case of a deficit of one or more nutrients
3. During a stressful situation of any kind

### Effects:

- allows better rooting
- improves root hair formation, root length and thus nutrient intake from the soil
- prolongs the period of vegetation of fruit vegetables and fruits
- improves metabolism and natural resistance to diseases and pests
- increases the overall vitality of plants
- Helps increase soil fertility by supporting soil microflora (revitalizes the soil) - this means that the fertilizer used is not only retained on the leaf, but a certain part always reaches the surface and thus, act as food for soil microflora
- prevents the accumulation of heavy metals and harmful substances in plants.

### Contents:

	Rokohumin Classic	Rokohumin
Total nitrogen (N)	4 % in dry matter	14 % in dry matter
phosphorus (P <sub>2</sub> O <sub>5</sub> )	9 % in dry matter	9 % in dry matter
potassium (K <sub>2</sub> O)	14 % in dry matter	14 % in dry matter

it further comprises amino acids, humic acids, S, B, Fe, Zn, Cu, Mn, Mo.

Amino acids in a fertilizer are needed for plant growth, regeneration and development. They are also of great importance for soil microflora by promoting the activity of soil microorganisms and enzyme activity. Plants are able to synthesize all the amino acids they need. Many amino acids are precursors or activators of phytohormones and growth agents in plants. In stressful situations, a plant can accumulate a significant amount of free amino acids that serve as a protective mechanism. The importance of this accumulation is that the plant spends less energy on protein synthesis. If the amino acid is supplied in the form of foliar nutrition, the plants will be in better conditions, which will affect growth and development. Thus, when a plant has individual building elements available and does not have to create them, its structure is growing at a faster rate.



Scan QR code for more information

### Effects of a few amino acids

**PROLINE** – contributes to the strength of the cell wall, increases the resistance of plants to stress factors and reduces the risk of damage, increases the ability of the pollen to fertilize the plant and the fruit seed.

**GLICINE** – Increases chlorophyll concentration and improves photosynthetic process conditions, has a positive effect on pollination and fruit formation.

**GLUTAMIC ACID** – affects osmotic processes in protoplasm and influences the opening and closing of vents, has a positive effect on pollination, activates seed germination, participates in nitrogen metabolism and protein synthesis.

**METHIONINE** – is a precursor of ethylene, enhances root development, regulates the opening of vents.

**TRYPTOPHAN** – an auxin precursor (a phytohormone that promotes the growth and strengthening of young roots, stimulates the growth of meristematic tissues), helps to overcome stress and prevents growth retardation.

**ARGININE** – increases the synthesis of hormones associated with the formation of flowers and fruits and facilitates the penetration of soil nutrients into the roots.

**ALANINE, VALINE and LEUCINE** – help to improve the quality of fruits.

**HISTIDINE** – promotes germination.

Nutrients from the leaf surface are most rapidly absorbed and maximally utilized by the plant. They improve leaf quality parameters in nutrient uptake, photosynthesis and other biochemical processes (leaf area size, chlorophyll content).

When plants cannot receive nutrients from the soil?



- Insect damage
- Lack of moisture
- Unsuitable soils pH
- Low temperature
- High temperature



# LOOSE BASIC FERTILIZER



**ROKOHUMÍN LOOSE is a universal organic fertilizer, suitable for every gardener and grower of vegetables, small fruits, vines, fruit trees, houseplants, ornamental shrubs, garden flowers and lawns (petunias, geraniums, orchids, ferns, bonsai, etc.).**

## Effects of Rokohumin Loose:

- It is suitable for all crops that are sensitive to burning or fertilization with nitrogen.
- Binds toxic metals in the soil and thus prevent their accumulation in plants.
- Promotes germination, rooting and plant viability and pest resistance.
- Enables better intake of macro (N, P, K) and micronutrients (Fe, Zn and others) by the root system and thus prevent the occurrence of plant chlorosis.
- Regulates the water regime in the soil, enriches the soil with microorganisms by providing them with optimal conditions and nutrients in the form of amino acids for their development and reproduction.
- Micro-organisms in the soil are repelled by the release of originally inaccessible nutrients and thus to better plant growth with less need for fertilizers.
- The fertilizer also contains small amounts of phosphorus in its natural form, as well as other nutrients and trace elements.
- The ecologically friendly form of gradually releasing organic nitrogen prevents the accumulation of harmful nitrates in plants and fruits of vegetables or small fruits, prevents the pollution of groundwater with unwanted nitrates.
- A special feature of the hooves and horns is their ability to bind nitrogen from the air.
- Versatility and a markedly ecological character are the main advantages of this fertilizer

## Contents:

- min. 7% N in dry matter, organically bound nitrogen
- min. 49.0% humic acids in dry matter

## Application rate:

With the fertilizer Rokohumín, we fertilize during the entire growing season, as well as during soil preparation before sowing or planting. We will carry out the application by even sowing and its subsequent incorporation into the soil. At a medium dose of 60 g / m<sup>2</sup> (1.5 kg per 25 m<sup>2</sup>), this use appears to be optimal. On poorer soils, a minimum of 80 g / m<sup>2</sup> (2.0 kg per 25 m<sup>2</sup>) is recommended. For houseplants 1-2 teaspoons (about 25 g) and then work into the soil. Add 1-2 teaspoons of fertilizer to the soil in the hole under the seedlings or, if it is already planted, apply around the seedlings. For ornamental plants, fruit trees, etc. apply 3 to 4 large spoons.

Thanks to the fertilizer, plant gains in strength and growth stability. The humic component, which directly affects the fertility of the soil, helps keep the water in the soil. It significantly helps to increase the chlorophyll content and the efficiency of photosynthetic function.



Scan QR code  
for more information

# GERMINATION ACTIVATOR



Is an stimulant excipient on an organic basis affecting the rooting and initial phase of plant growth.

#### Advantages:

- stimulates plant growth and development
- returns the dressed seed to its original physiologically active state
- due to several components of Rokoaktiv, the root system is intensively formed and developed and thus intensive intake of nutrients (macro- and microelements) and water is intensified, making the crop optimally prepared for wintering in the case of winter crops. In the case of other crops, like spring plants, it helps with better start and durability.
- the excellent condition of the crop helps to cope with less favourable conditions

#### Effects:

- Supports the formation of fine root capillaries. As a result, it increases the use of moisture and nutrition.
- Stimulates growth and yield. In particular, it promotes the flow of metabolites into seeds and fruits.
- Increases seed size. Especially with a good supply of nutrients.
- Favourably affects the N content of food wheat grain.

It increases germination energy and significantly affects germination speed and quality. It allows balanced emergence of stands and increases photosynthesis of germinating plants.

#### The effect of Rokoaktiv is exhibited by:

- a) better seed emergence
- b) promoting vegetative growth,
- c) accelerating the development of the whole plants,
- d) changing and optimizing the root-above roots ratio,
- e) increased yield and, in most cases, higher nutrient intake and improved plant health

**The aim is to return the seed to its original biological active state, that is, as it had before the seed had been dressed.**

Rokoaktiv is widely used as a root for cuttings or young seedlings. It significantly contributes to the creation, development and regeneration of the root system and above-ground parts of flowers and decorative trees.

#### Contents:

Total nitrogen (N)	5 % in dry matter
phosphorus (P <sub>2</sub> O <sub>5</sub> )	9 % in dry matter
potassium (K <sub>2</sub> O)	23 % in dry matter
Humic acids	20 % in dry matter
Dry matter content	20 %

it further comprises amino acids, S, B, Fe, Zn, Cu, Mn, Mo.

#### Application and dosage:

- added during seed dressing / tank-mix (TM)
- a dose of 4 - 6 litres per 1 ton of seed

SEED DRESSER  
+ ROKOAKTIV



SEED DRESSER  
+ other, rival stimulator



SEED DRESSER



Scan QR code  
for more information

# SOIL ACTIVATOR



Rokolan is an amino acid fertilizer intended for incorporation into the soil before sowing.

### Effects of Rokolan:

- Activates the soil microflora by providing sufficient nutrients to the microorganisms in the soil and thus, directly it contributes to a significant increase in their total quantity and to the revitalization and regeneration of the soil
- Indirectly helps to make nutrients in the soil available, by increasing the number of microorganisms in the soil, it catalyses their processes in the soil therefore making more nutrients available to the plant root system
- Reduces plant demand on the amount of mineral fertilizers used
- Helps to break down digestible residues after sowing by supporting the process by intensifying the activity of microorganisms

The nitrogen cycle in nature consists of three main processes that are conditioned by the action of microorganisms:

- Synthetic processes that produce organic nitrogen substances (the binding of atmospheric nitrogen by nitrogenic bacteria),
- Degradation processes by which the mineralization of organic nitrogen substances is carried out (ammonization),
- Conversion of mineral nitrogen compounds (nitrification, denitrification).

### Contents:


Nitrogen in organic form (N) in dry matter	min. 4,5%
Phosphorus (P <sub>2</sub> O <sub>5</sub> ) in dry matter	min. 13,0%
Potassium (K <sub>2</sub> O) in dry matter	min. 19,5%
Dry matter v %	min. 30,0%
Combustible substances in dry matter	min. 50,0%
<b>Amino acids</b>	<b>18</b>

### Dosage:

recommended dose per hectare: 50 liters / ha



Drill applying a liquid fertilizer



Application of Rokolan to the soil

*Doran and Parkin (1994) defined soil quality as "the capacity of a soil to function within ecosystem boundaries to sustain biological productivity, maintain environmental quality and promote plant and animal health"*



Scan QR code for more information

# APPLICATION TO VARIOUS CROPS

**DENSELY  
SOWN  
GRAIN**



15

**RICE**



16

**COTTON**



17

**SUNFLOWER,  
SESAME**



18

**RAPESEED**



19

**MAIZE**



20

**LEGUMINOUS  
PLANTS**



21

**VEGETABLES**



22

**BULB  
VEGETABLES**



23

**ROOT  
VEGETABLES**



24

**FRUITS**



25

**VINES**



26

**GRASSLAND**



27

**POTATOES**



28


**SUGAR  
BEET**



29

# DENSELY SOWN GRAIN

Crop	Repeat	Growth phase	Dose per ha
densely sown grain, sorghum, millet	2-3x with protection against diseases/pests	1. End of tillering 2. Stable 3. Post- flowering (mild maturation phasei – food wheat)	5 litres of Rokohumin / 150L of water



**Accompanying effect** - improving crop condition, optimizing root system growth, optimizing nitrogen uptake.

**Achieved effect** - improves the quantitative and qualitative parameters of grain.

**Tab. The effect of ROKOHUMIN on the structural parameters and yield of winter wheat**

No.	Treatment	Yield t/ha
1	Inspection ( without fertilizers)	4,28
2	<b>N30P45K45+ROKOHUMIN – 2x5L/ha</b>	<b>6,52</b>
3	N <sub>30</sub> P <sub>45</sub> K <sub>45</sub>	6,28



ROKOHUMIN treatment of winter wheat was carried out twice during the growing season - in a full seedling stage and before the first flag leaf appeared - a single dose - 5l / ha.

Research results have shown that ROKOHUMIN improves crop quality indicators and winter wheat yield levels.



# RICE

Crop	Repeat	Growth phase	Dose per ha
rice	1-3 x with protection against diseases/pests	1. Stage of first leaves. 2. Active tillering phase 3. Immediately after the start of the late reproductive phase, just before flowering	2x5 litres of Rokohumin / 200l of water




**Accompanying effect** - optimization of root formation, good vegetation condition, maintenance of vegetation in case of lack of moisture or in excess of moisture, stimulates active consumption of plant nutrients from the soil, reduces the temperature level, moisture and chemical stress, stimulates the growth of vegetative mass.

**Achieved effect** - increases yield up to 40%, accelerates cotton ripening, which makes it possible to harvest 75% of the crop before the rains, get the bulk of the raw cotton without yellowness, Increases the mass of 1000 seeds up to 10%, which allows you to get an additional premium for the harvest. provides improved fibre and seed quality, seed formation with higher sowing qualities. The treatment of fields with Rokohumin improves the structure of the soil and contributes to the accumulation of humus in it. Water and air permeability of the soil improves. Increasing the size, weight of rice grains and the quality of the crop.



# COTTON

Crop	Repeat	Growth phase	Dose per ha
cotton	1-2 x with protection against diseases/pests	1. Stage of 3-4 leaves 2. Budding phase of growth	2x5 litres of Rokohumin / 150l of water or 10 litres of Rokohumin / 200l of water



**Accompanying effect** - optimization of root formation, good vegetation condition, maintenance of vegetation in case of lack of moisture or in excess of moisture, stimulates active consumption of plant nutrients from the soil, reduces the temperature level, moisture and chemical stress, stimulates the growth of vegetative mass.

**Achieved effect** - increases yield up to 40%, accelerates cotton ripening, which makes it possible to harvest 75% of the crop before the rains, get the bulk of the raw cotton without yellowness, Increases the mass of 1000 seeds up to 10%, which allows you to get an additional premium for the harvest. provides improved fibre and seed quality, seed formation with higher sowing qualities. The treatment of fields with Rokohumin improves the structure of the soil and contributes to the accumulation of humus in it. Water and air permeability of the soil improves.



# SUNFLOWER, SESAME

Crop	Repeat	Growth phase	Dose per ha
sunflower, sesame	1-2 x with protection against diseases/pests	1. Stage of 6-8 leaves 2. Height of stand approx. 1 m (never in blossom phase!)	2x5 litres of Rokohumin / 150l of water or 10 litres of Rokohumin / 200l of water



**Accompanying effect** - optimization of root formation, good vegetation condition, support of inflorescence deployment, maintenance of vegetation in case of lack of moisture or in excess of moisture, optimization of nitrogen uptake.

**Achieved effect** - increases the number of grains and thus the overall crop, increases oil content.



**Tab. Sunflower harvest in t.ha<sup>-1</sup> at standard humidity**

Variant	2018	2019
<b>Rokolan</b> - 50l/ha per 150l of water (before sowing)		
<b>Rokoaktiv</b> - 5 l/t (seed dressing)		
<b>Rokohumin</b> - 2x 5l per 150l of water (foliar application)	3,83	3,75
<b>N<sub>15</sub>P<sub>15</sub>K<sub>15</sub></b> - 200kg/ha (sowing under heel)	2,37	3,42
<b>Non-fertilized control</b>	1,48	1,66

Foliar application 2x5l:  
1st application in phase of 6-8 leaves  
2nd application at height of stand approx. 1 meter (not during blossoming)  
or 1x10l per 200l of water


Despite the dry weather in 2018 compared to 2019, the following fact was confirmed. Seed treated with ROKOAKTIV had a better start of germination and early growth than untreated seed. The emergence of the treated stand was 7-10 days earlier than the untreated stand. Untreated vegetation emerged after rainfall. Longer vegetation time and foliage application by ROKOHUMIN resulted in the crops produced.



The use of industrial fertilizers leads to soil degradation, causing a loss of microflora in the soil. Rokolan is a way to improve soil quality and increase soil humus.

# RAPESEED

Crop	Repeat	Growth phase	Dose per ha
rapeseed	2-3 x with protection against diseases/pests	1. Spring during leaf covering min. 40-50% of soil 2. In the butonization stage 3. Before flowering	5 litres of Rokohumin / 150l water



**Accompanying effect** - improving the condition of the stand, optimizing the growth of the root system, optimizing nitrogen uptake, increasing resistance to cold and drought, strengthening the stand.

**Achieved effect** - maintaining HTS at lack of moisture, increasing HTS at normal humidity.


**Tab. Impact of ROKOHUMIN on rapeseed yield**

Variant	Yield t.ha <sup>-1</sup>
control	3,15
Rokohumin 3 x 5l/ha	3,52



# MAIZE

Crop	Repeat	Growth phase	Dose per ha
maize	1-2 x with protection against diseases/pests	1. 6-8 leaf stage 2. Height of stand approx. 1 m (never in blossom!)	2x5 litres of Rokohumin / 150 litres of water or 10 litres of Rokohumin / 200l water



**Accompanying effect** - optimization of root formation, good condition of the stand, support of inflorescence, maintenance of stands in case of lack of moisture or in case of excess moisture, optimization of nitrogen uptake.

**Achieved effect** - increase in the number of grains and HTS and thus in the overall harvest.



1 Rokohumin  
2, 3, 4 different fertilizers



**Tab. Maize yield in t.ha<sup>-1</sup> at standard humidity**

Variant	2018	2019
<b>Rokolan</b> - 50l/ha per 150l water (before sowing)		
<b>Rokoaktiv</b> - 5 l/t (seed dressing)		
<b>Rokohumin</b> - 2x 5l na 150l vody (foliar application)	7,94	10,09
<b>N<sub>15</sub>P<sub>15</sub>K<sub>15</sub></b> - 200kg/ha (sowing under the heel)	7,56	10,21
<b>Unfertilised control</b>	5,67	7,24

Foliar application 2x5l:

1st application in stage of 6-8 leaves

2nd application at height of stand 1m (not during flowering)


or 1x10l per 200l water

Despite the dry weather in 2018 compared to 2019, the following fact was confirmed. Seed treated with ROKOAKTIV had a better start of germination and initial growth phase than untreated seed. The emergence of the treated stand was 7-10 days earlier than the untreated stand. Untreated stands did not appear until after the rainfall. Longer growing time and leaf application with ROKOHUMIN was reflected in the resulting crops.

As a result of the use of industrial fertilizers, soil degradation occurs, which causes a loss of microflora in the soil. Rokolan is a way to improve the quality properties of the soil and increase humus in the soil.

# LEGUMINOUS PLANTS

Crop	Repeat	Growth phase	Dose per ha
leguminous plants	2-3 x with protection against diseases/pests	1. Stage of young plants till 7 leaves 2. Period from main growth till start of flowering 3. flowering period – Teat formation	5 litres of Rokohumin / 150-200l of water



**Accompanying effect** - higher establishment of the lower floor of the branches, increased flow of assimilates into the seeds, increased drought resistance, support of root system formation, improvement of the condition of the stand.

**Achieved effect** - increase of nitrogen intake, maintenance of HTS in case of lack of moisture, increase of HTS at normal humidity, increase of yield.

**Tab. Impact of Rokohumin on soy yield (Taken into account with 14% humidity)**

Variant	Yield t.ha <sup>-1</sup>
Control	2,78
Rokohumin 2 x 5l	3,18



# VEGETABLES

Crop	Repeat	Growth phase	Dose per ha
vegetables	every 10-14 days	during the period of main growth (it is recommended to use a wetting agent)	5 litres of Rokohumin / 200l water



**Tab. 1 Impact of Rokohumin on tomato yield**

Fertilizer	Yield			Yield of commodity production			Marketability %
	growth to standard		%	growth to standard		%	
	t/ha	t/ha		t/ha	t/ha		
Non-fertilised control	39,7	-	-	36,5	-	-	91,9
NPK+Rokohumin	46,3	6,6	16,6	43,4	6,9	18,9	93,7



**Tab. 2 Impact of Rokohumin on cucumber yield**

Fertilizer	Yield			Yield of commodity production			Marketability %
	growth to standard		%	growth to standard		%	
	t/ha	t/ha		t/ha	t/ha		
control	12,95	-	-	11,57	-	-	89,3
Rokohumin (foliar application in 3rd period)	15,02	2,07	16	13,66	2,09	18,1	90,9
NPK	16,25	-	-	14,58	-	-	89,7
NPK+ Rokohumin (foliar application)	18,82	2,57	15,8	16,9	2,32	15,9	89,8




**Accompanying effect** - intensive formation of the root system, higher resistance to weather fluctuations.

**Achieved effect** - increasing nitrogen intake, creating conditions for higher and better yields.




# BULB VEGETABLES

Crop	Repeat	Growth phase	Dose per ha
bulb vegetables	2 x after creating a sufficient leaf area	during main growth stage (on onion, garlic and chives it is essential to use wetting agent)	5 litres of Rokohumin / 200l water



Tab. Impact of Rokohumin on onion yield

Variant	Yield t/ha
Control (without fertilisers)	9,88
NPK (local - standard)	10,2
NPK (local) + Rokohumin 5l/ha	11,34



**Recommendation for better yield and protection against diseases and pests for garlic:**

Before planting the clove of garlic, soak in a 10% solution of Rokoaktiv with water in a ratio of 1:10 for 12 hours. Fertilize according to the application table.


**Accompanying effect** - intensive formation of the root system, higher resistance to weather fluctuations.

**Achieved effect** - increasing nitrogen intake, creating conditions for higher and better yields.



# ROOT VEGETABLES

Crop	Repeat	Growth phase	Dose per ha
root vegetables	2-3 x according to the condition of the stand and the course of weather/wind conditions	During spring, leaf covering of soil by plants min. 40-50% 2-3. Two-three-week intervals	5 litres of Rokohumin / 150l water



**Accompanying effect** - optimal condition of the stand, enlargement of the bougainvillea, root.

**Achieved effect** - increasing nitrogen intake and its effective use, creating preconditions for higher and better yields.

**Tab. Impact of Rokohumin on carrot yield**


Fertilizer	Yield			Yield of commodity production			Marketability %
	t/ha	t/ha	%	t/ha	t/ha	%	
Non-fertilised control	32,9	-	-	29,3	-	-	89,1
NPK+Rokohumin	37,4	4,5	13,7	34,4	5,1	17,4	92

Rokohumin also has a positive effect on the biochemical parameters of carrots by increasing the content of carotene, ascorbic acid and reducing the content of nitrates.



# FRUITS

Crop	Repeat	Growth phase	Dose per ha
fruits	3-4 x with protection against diseases/pests	1. Young growing twigs 2. 14-21 days after 1st application 3. 2-3 weeks after 2nd application	5-10 litres of Rokohumin / water as needed



**Accompanying effect** – increase of chlorophyll content in leaves, increase of number of leaves, higher flow of assimilates into fruits, significantly improvement of health condition.

**Achieved effect** - better condition of trees, increased sugar content, larger fruits.

**Clients’ feedback when using Rokohumin foliar fertilizer for fruit orchards:**

When using Rokohumin fertilizer, they achieved a more even size and quality of fruit, thus achieving a higher % of marketable goods for direct consumption. The fruits have a more pronounced taste and aroma, there is an increase in sugar content and dry matter, which has a positive effect on storage. By applying the fertilizer, a larger number of pollinated flowers is preserved, which contributes to an increase in yield by 15% or more.



# VINES

Crop	Repeat	Growth phase	Dose per ha
vines	4-5 x with protection against diseases/pests	1. Application on young growing bine 2. Application after flowering 3. Subsequently in 2-3 weeks Intervals in combination with protection against diseases and pests	5-7 litres of Rokohumin / water as needed



**Accompanying effect** - increased flow of assimilates into berries, immediate effect on the biosynthesis of sugars and acids.

**Achieved effect** – increasing sugar content and extract with a slight increase in yield, a significant improvement in health strengthening the growth and creating the condition for increasing the harvest.



ROKOAktiv+ROKOHUMIN



Unfertilised

**Experiences:**

When planting a vineyard and achieving a healthy and strong growth, we recommend soaking the roots in a solution of Rokoaktiv with water in a ratio of 1:10 for 12 hours before planting. For watering, we recommend Rokohumin at a dose of 2-3 times at 35L / ha or use the rest of Rokoaktiv. With an existing vineyard, we fertilize according to the application table.

**Client feedback:**

- resistant growth during the whole vegetation period against diseases, pests
- maturation of wood mass after harvest
- significant increased yield per ha
- increasing the concentration of aromatic substances transferred to the wine
- faster acquisition of sugar content required at harvest, this affects the earlier harvest

# GRASSLAND

Crop	Repeat	Growth phase	Dose per ha
grassland	4 x	1. At the beginning of vegetation after creating sufficient leaf area at stand height of 10-15 cm 2-4. After regeneration, after mowing at a stand height of 10-15cm	5 litres of Rokohumin / 150l water




**Accompanying effect** - high nutritional value of grasslands, stimulation of the root system, increasing the density of all grasslands.

**Achieved effect** - earlier time of obtaining commercial biomass, quantitative increase, increase of leaf biomass, high feed value of obtained production.

# POTATOES

Crop	Repeat	Growth phase	Dose per ha
potatoes, cassava, yam, sweet potatoes	3 x with protection against diseases/pests	1. During spring, leaves covering the soil min.40-50% 2. a 3. If the need for application of agents protecting the plants arises	10 litres of Rokohumin / 200l water



**Accompanying effect** - support of root system formation, support of stand growth, increase of the number of medium and smaller tubers in seed potatoes, increase of starch content in industrial potatoes, increase in the number of tubers in ware potatoes.

**Achieved effect** - increase of nitrogen intake and its effective use for crop production, optimal condition of plants, increase of chlorophyll concentration, enlargement of leaves and their slower aging.


**Tab. Impact of Rokohumin potato yield**

VARIANT	Yield t.ha <sup>-1</sup>
unfertilised	24,24
NPK	31,06
NPK+Rokohumin (2 x 5l/ha)	37,22



# SUGAR BEET

Crop	Repeat	Growth phase	Dose per ha
sugar beet	3-4 x with protection against diseases/pests	During spring, leaves covering the soil min 40-50% 2. a 3. In two-three-week intervals 4. Late summer (July)	5 litres of Rokohumin / 150l water



**Accompanying effect** - increase of nitrogen intake and its effective use, optimal condition of the stand, enlargement of bulbs, roots.

**Achieved effect** - late summer application (July) can increase the sugar content of the bulbs by up to 2%, higher and better yield.

**Tab. Impact of Rokohumin on sugar beet yield**

Variant	Yield t.ha <sup>-1</sup>	Sugar content%
Unfertilised	38,04	18,50
NPK	44,56	18,89
NPK+Rokohumin (3 x 5l)	51,24	19,68



## Contents:

Total nitrogen (N)	5 % in dry matter
phosphorus (P <sub>2</sub> O <sub>5</sub> )	9 % in dry matter
potassium (K <sub>2</sub> O)	23 % in dry matter
humic acid	20 % in dry matter
<b>amino acids</b>	<b>18 types</b>
dry matter content	20 %
S, B, Fe, Zn, Cu, Mn, Mo.	

Rokoaktiv is an organic-based stimulant adjuvant acting as a germination activator with a stimulating effect on rooting and the initial stage of plant growth. Rokoaktiv prevents the negative effect of the feed dresser, increases the percentage of seed germination and ensures a better start in the initial phase of growth due to better root system development and a starting dose of nutrients with a high content of amino acids and humic substances.




The seed is inhibited by germination, growth and development due to the use of the feed dresser, which includes an effective slat. The goal of Rokoaktiv is to return the physiology of the plant to its original state. In the experiments, we closely monitored and

recorded the factors and influences on changes in germination, rooting, growth intensity, the ratio of roots to the aboveground part and the strength of the cell structure, along with the balance of growth.



Germination of seeds in relation to seed dressing and dose of activator of germination - Rokoaktiv



In 2018 and 2019, we carried out an experiment with the Rokoaktiv stimulator at the experimental workplace of the Agroecology Research Institute in Michalovce. The experiments at the workplace in Milhostov were based on two crops, sown maize and annual sunflower.



The emergence of vegetation is significantly influenced by the weather, which means that the seed treated with the Rokoaktiv stimulator emerged 10 days earlier than untreated. These did not arise until after the rainfall. After generalization, it can be emphasized that in the dry and warm weather, as in 2018, in the experiment with sown maize and annual sunflower, the highest yields were achieved in the treatment of seeds with Rokoaktiv. Higher resistance to stress conditions, drought was shown.

### Rokoaktiv in the conditions of horticulture, viticulture and cultivation of decorative trees and flowers-roots

Rokoaktiv is widely used as a rooter for cuttings or young seedlings and as a regenerating agent for damaged roots of larger plants or woody plants.

The unique composition of Rokoaktiv, with its stimulating effect, contributes to the formation and growth of the plant root system.

The effect of Rokoaktiv in the conditions of fruit growers, winemakers, growers of decorative trees and flowers:

- best tree acceptance
- best condition of the stand
- faster onset of the fertility phase of fruit trees and vines
- for flowers and decorative trees, rapid regeneration of the root system and the aboveground part of the plants

Before planting, soak the roots in Rokoaktiv with water in a ratio of 1:10 for 12 hours. We will use the

rest of Rokoaktiv for watering. Additional fertilization to the roots of already planted trees and vine sprouts is applied using a pruner or hydraulic drill.



## Effects of Rokoaktiv as an activator in biological and efficiency in the economic field of agriculture in winter wheat cultivation.

### 1. Impact of Rokoaktiv activator on winter wheat grain quality indicators:

Variant	Crude protein, % in dry matter	Crude protein yield, kg/ha	Yield q/ha	Pp, g/kg feed unitz yield	Security 1 k.j. Pp, g	Yield KJP, q/ha
Unfertilised (seed dressed by Polaris, ME, 1,5l/1t seeds)	13,7	401	41,1	106	88	42,1
N <sub>80</sub> +40+30P <sub>15</sub> K <sub>30</sub> Background (seed dressed by Polaris Polaris, ME, 1,5l/1t seeds)	13,6	527	54,7	105	87	55,6
<b>Background+Rokoaktiv, seed treatment before sowing, 5l/t seeds</b>	<b>13,8</b>	<b>564</b>	<b>57,4</b>	<b>107</b>	<b>89</b>	<b>58,9</b>

The use of the fertilizer Rokoaktiv contributed to the increase of the yield of crude protein by 37 kg / ha, feed and feed protein units by 2.7 and 3.3 q / ha, or by 6% on average compared to the mineral background N<sub>80</sub>+40P<sub>25</sub>K<sub>35</sub>.

### 2. Impact of activator Rokoaktiv on the total grain yield of winter wheat.

Variant	Total Yield q/ha	+/-, q/ha		Dry matter Yield q/ha
		To control	To background	
Unfertilised (seed dressed by Polaris, ME, 1,5l/1t seeds)	34.0	-	-	29,2
N <sub>80</sub> +40+30P <sub>15</sub> K <sub>30</sub> -Background (seeds dressed by Polaris, ME, 1,5l/1t seeds)	45.2	11,2	-	38,9
<b>Background+Rokoaktiv, seed treatment before sowing, 5l/t seeds</b>	<b>47.4</b>	<b>13,4</b>	<b>2,2</b>	<b>40,8</b>

Pre-sowing treatment of winter wheat seeds with Rokoaktiv activator had a positive effect on the total yield of winter wheat grain - a statistically reliable increase of crop grain of 2.2 q / ha was achieved at a yield of 47.4 q / ha, which is 5% more compared to the background. In the experiment, a maximum dry matter yield of 40.8 q / ha was achieved.

Related to the pre-sowing treatment of winter wheat seeds with Rokoaktiv activator at a dose of 5 liters per 1 ton of seeds in combination with seed dresser in growing the culture, high efficiency was confirmed at the minimum financial costs of using 1 liter of Rokoaktiv for seeds needed per 1-hectare areas of winter wheat.



## Contents:

Nitrogen in organic form(N) in dry matter	min. 4,5%
phosphorus (P <sub>2</sub> O <sub>5</sub> ) in dry matter	min. 13,0%
potassium (K <sub>2</sub> O) in dry matter	min. 19,5%
dry matter in %	min. 30,0%
combustible substances in dry matter	min. 50,0%
<b>amino acids</b>	<b>18 types</b>

Rokolan is a basic fertilizer on an organic basis intended for root nutrition of agricultural crops. Its effect provides a better start in the initial phase of growth due to better development of the root system and the starting a dose of nutrients with a high amino acid content. Rokolan is a substitute for manure.

Due to the use of industrial fertilizers, water and wind erosion, soil degradation occurs, ie a reduction in the content of humic substances, a reduction in the formation of microbial biomass and an unnatural reduction in biological activity in the soil.

Rokolan is a way to improve the quality properties of the soil and increase the amount of microorganisms in the soil - to increase humus in the soil.

Rokolan application before sowing



Experiment in annual sunflower - Rokolan at a dose of 50l / ha



Experiment on sown maize - Rokolan at a dose of 50l / ha



Tab. Sunflower harvest in t.ha<sup>-1</sup> at standard humidity

Variant	2018	2019
<b>Rokolan</b> - 50l/ha per 150l of water (before sowing)		
<b>Rokoaktiv</b> - 5 l/t (seed dressing)		
<b>Rokohumin</b> - 2x 5l per 150l of water (foliar application)	3,83	3,75
<b>N<sub>15</sub>P<sub>15</sub>K<sub>15</sub></b> - 200kg/ha (sowing under the heel)	2,37	3,42
<b>Unfertilised control</b>	1,48	1,66

Leaf application 2 x 5l:

1st application in the stage of 6-8 leaves

2nd application at a stand height of 1 meter (not at flowering)

or 1 x 10l per 200l of water

Tab. Maize yield in t.ha<sup>-1</sup> at standard humidity

Variant	2018	2019
<b>Rokolan</b> - 50l/ha per 150l of water (before sowing)		
<b>Rokoaktiv</b> - 5 l/t (seed dressing)		
<b>Rokohumin</b> - 2x 5l per 150l of water (foliar application)	7,94	10,09
<b>N<sub>15</sub>P<sub>15</sub>K<sub>15</sub></b> - 200kg/ha (sowing under the heel)	7,56	10,21
<b>Unfertilised control</b>	5,67	7,24

Leaf application 2 x 5l:

1st application in the stage of 6-8 leaves

2nd application at a stand height of 1 meter (not at flowering)

or 1 x 10l per 200l of water

# ROKOLAN



Seed drill with liquid fertilizer applicator

After application of Rokolan to the soil and subsequent activation of the microflora, which releases bound nutrients in the soil, the plants had a guaranteed supply of otherwise unacceptable nutrients. The more the plant drained, the more microorganisms replenished. Even after the nutrients were depleted by the plants, a good to very good supply of these nutrients remained in the soil. In the experiment of sown maize and annual sunflower, the highest yields were achieved with technologies such as the activator Rokoaktiv, application of Rokolan before sowing and leaf application of Rokohumin. Rokoaktiv acts to support seed germination and the initial phase of growth. Rokolan, in turn, acts as a support for the microflora in the soil, which makes nutrients available in the soil. Rokohumin then directly supports the plant and supplies the necessary nutrients.



## Fertilization of orchards and vineyards with liquid amino acid fertilizer Rokolan

Fertilization of the soil with the vineyard / orchard already established by Rokolan is applied with a coulter to a depth of 25-30 cm, solitaires with a hydraulic drill or watering.



Rokolan application after planting

## Rokolan - technical crops

Technical crops intended for canning are fertilized and irrigated by drip irrigation in the presence of amino acid fertilizer Rokolan.

### Advantages of drip irrigation:

It is ideal to supply nutrients to the plant in several stages (4x15l Rokolan per hectare), using Rokolan diluted with water. The essence is to distribute nutrients to the plants evenly throughout the growing season.



Drip irrigation



Hydro drill



# INNOVATIONS

# IN SUPPORT OF FULFILLING THE GENETIC POTENTIAL OF A PLANT

## Use of drones in agriculture

### 1. MONITORING

### 2. PLANT NUTRITION

The importance in applying the liquid fertilizer Rokohumin by drones is in lower and more frequent doses. This way, the plant can get closer to the genetic potential of its fertility. The advantage is the possibility of applying liquid fertilizers during the entire growing season and to all the plants.





# INTERNATIONAL ACTIVITIES



**NIGERIA**





**NIGER**






# TANZANIA



# KENYA





# EQUATORIAL GUINEA





**RWANDA**



**ETHIOPIA**





配方肥称量



配方喷施

CHINA



第三次配方肥喷施



配方喷施



罗克歌微生态氨基酸配方肥在番茄上的应用效果研究

**试验目的:** 受新疆阿克苏“罗克歌有限公司 (ROKOSAN s.r.o.)”委托,对其研发的微生态氨基酸配方肥进行应用效果研究,通过测量番茄的物候期、坐果量、干物质、产量、品质、商品率、抗病性等主要指标,分析得到产品应用效果,以期找到最佳配方组合,为推广应用提供理论依据。

**试验材料:** 供试配方:氨基酸叶面肥-复合微生物菌剂  
供试品种:改良金满9号

**试验方法:** 随机区组,3个处理,3次重复,每个处理3槽,小区面积48.45平方米,株距20厘米,行距190厘米

**定植时间:** 2021年10月21日







# PILLARS OF A QUALITY YIELD



**ROK**  **AKTIV**

---

**Germination  
activator**

+

**ROK**  **LAN**

---

**Soil  
activator**

+

**ROK**  **HUMIN**

---

**Leaf  
nutrition**

# FOR HEALTHIER AND SELF-SUFFICIENT WORLD

**ROKOVIA**

**ROKOVIA**  
Krmanova 14  
Košice  
Slovak Republic

[www.rokovia.net](http://www.rokovia.net)

**ROKOSAN**

**ROKOSAN**  
Kollarova 446  
Sečovce  
Slovak Republic

[www.rokosan.com](http://www.rokosan.com)